



# Autologous and allogeneic transplantation of adipose derived stem cells have similar efficacy for type 1 diabetes mellitus therapy in mouse models

Anh Nguyen-Tu Bui<sup>1</sup>, Oanh Thi-Kieu Nguyen<sup>1</sup>, Cong Le-Thanh Nguyen<sup>1</sup>, Loan Thi-Tung Dang<sup>1</sup>, Dung Phuong Nguyen<sup>2</sup>, Phuc Van Pham<sup>1,3,\*</sup> , Ngoc Kim Phan<sup>1</sup>

<sup>1</sup>Laboratory of Stem Cell Research and Application, University of Science, Vietnam National University, Ho Chi Minh City, Viet Nam

<sup>2</sup>Faculty of Traditional Medicine, University of Medicine and Pharmacy, Ho Chi Minh City, Viet Nam

<sup>3</sup>Laboratory of Cancer Research, University of Science, Vietnam National University, Ho Chi Minh City, Viet Nam

## \*For correspondence:

pvphuc@hcmuns.edu.vn

**Competing interests:** The authors declare that no competing interests exist.

**Received:** 11 November 2016

**Accepted:** 10 December 2016

**Published:** 21 December 2016

Copyright The Author(s) 2016. This article is published with open access by BioMedPress (BMP).

This article is distributed under the terms of the Creative Commons Attribution License (CC-BY 4.0) which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

## Abstract

**Introduction:** Type 1 diabetes mellitus (T1D) disease is caused by lesions or dysfunction of beta cells of pancreatic islets, causing less insulin to be secreted into the blood and thereby increasing glucose levels in the blood. In this study, we evaluated and compared the efficiency of treatment for T1D using autograft and allograft adipose-derived stem cells (ADSCs). **Methods:** ADSCs were collected from the belly of mice before they were injected using a single dose of streptozotocin (100 mg/kg) to induce T1D. T1D mice were intravenously injected with a dose of  $2 \times 10^6$  ADSCs into the tail vein. Therapeutic efficacy was assessed by survival rate, blood glucose levels, serum insulin levels, histology and immunohistochemistry of pancreatic islets. **Results:** The results showed that both autograft and allograft transplantation of ADSCs demonstrated similarities in mortality rate, blood glucose level, blood insulin level, quantity and size of pancreatic islets. Both transplantations significantly improved T1D mice, which showed a decrease in mortality rate as well as blood glucose level, and increases in blood insulin level, quantity and size of pancreatic islets. **Conclusion:** The similar results suggest that both autologous and allogeneic transplantations of ADSCs are promising therapy for T1D treatment.

---

## Keywords

Adipose derived stem cells, type 1 diabetes mellitus, mouse model, autologous, allogeneic, autograft, allograft, stem cell transplantation

---

## Introduction

Diabetes mellitus disease is caused by lesions or dysfunction of beta cells in pancreatic islets, resulting in inadequate production and usage of insulin leading to abnormally high blood glucose levels. There are over 422 million diabetic patients in the world today and this number is expected to double by 2025 (Halban et al., 2014; Organization, 2016). Diabetes mellitus is a chronic disease with many complex symptoms related to blood glucose rise and associated ketoacidosis; such symptoms include slow healing cuts, blurred vision, numbness in feet and legs, and nausea and vomiting. There are two main kinds of diabetes mellitus: diabetes type 1 and type 2. The main general characteristic of both types is extremely high blood glucose level which cannot be controlled. Worldwide there are about 16 million type 1 diabetic patients, about 5% - 10% of all diabetic cases, mainly in children and adolescent (Halban et al., 2014; Organization, 2016).

Diabetes mellitus type 1 (T1D) patients often manifest other physiological characteristics, such as increased appetite and thirst, increased urination and weight loss (Organization, 2016). Although T1D can be treated by supplementing insulin into the body, a major disadvantage of this approach is the patients completely depend on exogenous insulin. The long-term usage of insulin may control blood glucose levels less and may cause lipodystrophy and other serious complications, such as retinopathy, blindness, cardiovascular disorders, stroke, kidney failure and cerebrovascular diseases.

Pancreas/islet transplantation has been one alternative therapy for diabetes treatment but currently stem cell transplantation has garnered wider interest. Pancreas/islet transplantation is challenging due to limited tissue source which impedes it from being a widespread application. New investigations into development of new therapies, such as stem cell therapy, for diabetes treatment have led to promising preclinical research results; stem cell transplantation have shown efficiency for diabetes treatment (Kraoz et al., 2013; Katuchova et al., 2015; Kwon et al., 2008; Lin et al., 2009; Yaochite et al., 2015).

Stem cells from fat tissue is an abundant source of mesenchymal stem cells (MSCs). Among the different kinds of MSCs, adipose derived stem cells (ADSCs) are a great stem cell candidate due to its abundant supply and resource. In fact,

the proliferation of ADSCs has been demonstrated to be better than cells from bone marrow tissue. Several research studies have demonstrated ADSCs are able to differentiate into multiple cell lines, including adipocytes, osteoblasts, chondrocytes and even insulin-producing cells of pancreatic islets (Bunnell et al., 2008; Karaoz et al., 2013). Moreover, ADSCs also secrete cytokines which are involved in immune modulation and which support metabolic regulation in the body. Due to these properties, ADSCs are considered as a promising candidate for stem cell transplantation for treating T1D (Lin et al., 2009; Lin et al., 2015; Paek et al., 2014; Zhou et al., 2016; Zuk, 2013).

Herein, we evaluated the use of ADSC transplantation to treat T1D; we investigated both autologous and allogeneic ADSCs. We hypothesized that the effects of autologous and allogeneic transplantations of ADSCs might be similar; both might be efficacious given the immune modulation property of ADSCs.

## Materials-Methods

### Cell isolation and proliferation

Murine ADSCs were isolated and proliferated according to a previously published protocol (Vu et al., 2015). Briefly, fat tissues were obtained from mice genital fat. Cell isolation was performed using ADSC Extraction kit (GeneWorld Ltd., Vietnam). After collection, cells were cultured in MSCCult medium (RegenMed Ltd., Ho Chi Minh, Vietnam) containing DMEM/F12 supplemented with 10 ng/mL epidermal growth factor (EGF), 10 ng/mL basic fibroblast growth factor (bFGF) and 10% fetal bovine serum (FBS), at 37°C and 5% CO<sub>2</sub>, and refreshed every 3 days; except for media all reagents were purchased from Sigma-Aldrich (St. Louis, MO).

### *In vitro* differentiation

ADSCs were induced to differentiate into adipogenic cells and osteogenic cells in specific differentiation medium. For adipogenic differentiation, cells were plated and cultured in DMEM/F12 supplemented with 0.5 µM dexamethasone, 50 µM indomethacin, 1.5 µg/mL insulin, and 0.5 µM 3-isobutyl-1-methylxanthine (IBMX) (all reagents were purchased from Sigma-Aldrich). After 5-7 days, the intracellular lipid accumulations were evaluated by Oil red O staining (Sigma-Aldrich, St Louis, MO). For osteogenic differentiation, cells were plated and cultured in DMEM/F12 medium supplemented with 10% FBS, 1% antibiotic-antimycotic 100X, 100 nM dexamethasone, 10 mM beta-glycerol, 10 µM ascorbic acid, 20.8 mg/mL L-leucine, and 100 ng/mL L-lysine (all purchased from Sigma-Aldrich). After 14-21 days, calcium accumulation in the extracellular matrix was confirmed by Alizarin Red staining (Sigma-Aldrich, St Louis, MO).

## **Inducing T1D mouse model by streptozotocin**

Mice weighing 25-28 g were selected and stably farmed for 2 weeks before induction of diabetes. All procedures and manipulations were performed in accordance with instructions and approval of the Animal Care Committee. The mice were fasted for 24 hours before streptozotocin (STZ) injection. STZ (purchased from Santa Cruz Biotechnology, Dallas, TX) was dissolved in cooled 0.01 M citrate buffer at pH 4.5. The T1D mice were induced by a single intraperitoneal injection of 100 mg/kg of STZ. Mouse blood glucose was examined by Accu-Chek Active (Mannheim, Germany). Unlike T1D mice which received STZ injection, control (normal) mice received PBS.

## **Mouse ADSC transplantation**

After 3 weeks, STZ-induced diabetic mice were intravenously injected with  $2 \times 10^6$  mouse ADSCs, either autologous or allogenic, in 200  $\mu$ L sterile phosphate buffered saline (PBS; Sigma-Aldrich) per mouse. The treated mice were monitored for survival rate, blood glucose levels, serum insulin and morphological analysis of pancreatic islets. The placebo treatment was injection with PBS only.

## **Glucose and insulin tolerance tests**

Glucose tolerance tests were conducted on the mice at 8 weeks after the transplantation. Mice were fasted 6-8 h before the glucose tolerance test. D-glucose was intraperitoneally injected into mice (1.5 g/kg body weight) and blood glucose levels were measured at 0, 15, 30, 60 and 120 minutes. Insulin was dissolved in PBS at a dose of 0.75 mg/kg body weight. Blood glucose levels of mice were measured at 0, 15, 30, 60 and 120 minutes after insulin treatment.

## **Serum insulin measurement**

At day 56, mouse blood was obtained from facial vein and serum was collected by centrifugation at 3000 rpm in 10 minutes at 4°C. The transparent faint yellow serum was assessed to measure insulin concentrations by Mercodia Ultrasensitive Mouse Insulin Enzyme linked immunosorbent assay (ELISA) (Uppsala, Sweden). The calibrator curve was constructed and insulin concentrations from the samples were extrapolated from the standard curve (analysis by Prism 6, GraphPad Software, San Diego, CA).

## **Histological analysis of pancreatic islets**

After 56 days of treatment, pancreases from mice were collected and fixed in 4% paraformaldehyde, treated with 30% sucrose overnight and embedded in optimal cutting temperature compound (OCT; Sigma-Aldrich). The 7  $\mu$ m-thin sections were stained with Hematoxylin & Eosin (Sigma-Aldrich) following the standard established procedure. The pancreatic islets were observed by

microscopy and were analyzed, counted and measured by Axio Vision Microscopy Software (Carl-Zeiss, Germany).

For immunofluorescence staining, the 7  $\mu\text{m}$ -thin sections were dipped in 1% bovine serum albumin (BSA; Sigma-Aldrich) for 1 hour at room temperature. The blocked slides were incubated with insulin antibody (Goat anti-Rabbit Alexa Fluor 488, Sigma-Aldrich) overnight at 4°C, then stained with FITC-conjugated secondary antibody (sc-2012, Santa Cruz Biotechnology) for 2 hours. Finally, the sections were stained with Hoechst 33442 (Sigma-Aldrich) for nucleus staining and observed by fluorescence microscopy (Cell Observer, Carl-Zeiss, Germany).


### Data analysis

The data were analyzed for statistical significance using GraphPad Prism software. Data were presented as mean  $\pm$  SD. When applicable, a Student's unpaired t-test and one-way ANOVA were used to determine significance;  $p < 0.05$  was considered to be statistically significant.

## Results


### Isolation and characterization of adipose derived stem cells

Genital fat of mice was collected according to the protocol of the ADSC Extraction kit. Mononuclear candidates exhibited a round shape (red arrow, **Fig. 1A**) and were a multiple mix of red blood cells and platelets. Red blood cells were less adhesive and discarded when refreshing culture media. After 24 h, mononuclear cells began to attach and spread along the culture surface. At day 3, all unattached cells and red blood cells were discarded (**Fig. 1B**) (Vu et al., 2015).



**Figure 1. Morphology of ADSCs in primary culture.** (A) after isolation, (B) after 3 days of culture, (C) after 5 days of culture, and (D) after 7 days of culture.

After 5 d, cells began to adhere and proliferate to about 30 – 50% of cultured areas. Cell morphology was not homologous; most cells had an elongated, fibroblast-like shape while some were shorter and flat, and some had a comma shape (**Fig. 1C**). From days 7-9, cell density reached about 70 – 80% confluence and cells could be sub-cultured for the 1<sup>st</sup> passage (**Fig. 1D**). The sub-culture procedure allows for more space and nutrition for cell proliferation; through each sub-culture non-ADSCs were removed to enable the ADSC population to become more homogenous.




**Figure 2. Cell morphology of mADSCs.** (A) Osteogenic cells on day 14, (B) staining of osteoblasts with Alizarin Red, (C) adipogenic cells on day 7, and (D) staining of adipocytes with Oil Red O.

Cell morphology changed when ADSCs were cultured in ‘differentiation’ medium (i.e. explain again what the medium was). After 7 -12 d in this medium, cells differentiated into adipocytes with lipid droplet formation and showed positive staining with Oil red O (**Fig. 2**). Similarly, for osteoblast differentiation, after 14 d ADSCs changed their shape, accumulated intracellular calcium ( $\text{Ca}^{2+}$ ) and magnesium ( $\text{Mg}^{2+}$ ), and showed positive staining with Alizarin Red (**Fig. 2**). ADSC candidates were capable of differentiating into adipocytes and osteoblasts *in vitro*. These results are similar to previous publications (Timper et al., 2006; Vu et al., 2015).

### ADSC transplantation reduced the mortality rate of diabetic mice

At 56 d after transplantation, except for the T1D-Placebo group, all other groups showed no mortality. In the control (normal) mouse groups which received autograft (autologous) or allograft (allogeneic) ADSC transplantation, the survival rate was 100%. Mice in these groups remained healthy, agile and active. These results showed that cell transplantation was safe for mice. Immediately after transplantation, mice did not get shocked or tired nor did they exhibit abnormal manifestation for the 56 days of follow-up. There were no differences between Normal-autograft group, Normal-allograft group and Normal-placebo group (mice which received placebo treatment) (Fig. 3).




**Figure 3.** Kaplan–Meier survival curve comparing between the groups after mADSC transplantation.

In T1D-placebo group, the health of the mice worsened and most mice became tired, lazy and unmotivated; only about 60% of mice survived at day 56 post transplantation. Due to the impact of STZ, the beta cells of pancreatic islets were destroyed, resulting in a high blood glucose level. Mice lost weight and exhibited multiple complications, such as corneal lesions, shuddering, and excitability. However, after cell transplantation, for both the T1D-autograft and T1D-allograft groups, the diabetic mice showed improvement of their health status; these included weight gain, smooth fur, and increased agility and activity. There were no mortality cases post cell transplantation, suggesting that stem cell transplantation could help prolong the life of T1D mice.

### ADSC transplantation reduced blood glucose level of diabetic mice


In the control (normal) mouse groups, the blood glucose level remained stable during the 56 d follow-up after treatment. There was no statistical difference between Normal-autograft group, Normal-allograft group or Normal-placebo group. The results showed that ADSC treatment (autograft or allograft), like placebo treatment, did not cause hyperglycemia in normal (non-diabetic) mice.



**Figure 4. Comparison of blood glucose levels (mg/dL) between groups after ADSC transplantation.**

However, in T1D diabetic mice, ADSCs affected the regulation of mouse blood glucose levels; the blood glucose levels of the autograft and allograft transplanted groups strongly decreased from day 7 to 28. At day 56 after transplantation, the blood glucose levels of the autograft and allograft groups were  $262 \pm 48$  mg/dL and  $233 \pm 36$  mg/dL, respectively, which was maintained from day 28 to day 56 (Fig. 4).

### Glucose and insulin tolerance tests



**Figure 5. Glucose (A) and insulin (B) tolerance tests.**




After D-glucose injection, all mice exhibited hyperglycemia. In T1D-Placebo group, blood glucose metabolism was slowed down such that after 120 minutes, blood glucose levels did not return as pre-tolerance. Blood glucose levels of the remaining groups recovered after 120 minutes (Fig. 5A). These results show that the treatment groups (T1D-Autograft and T1D-Allograft) have the ability to regulate blood glucose.

For insulin tolerance, serum insulin levels of all groups were reduced after insulin injection. Mice responded fairly to exogenous insulin and did not occur insulin-resistance after cell transplantation (Fig. 5B).


### Histological analysis of pancreatic islets and serum insulin measurements

After treatment, the pancreatic islets of T1D mice were recovered. The number of islets in the treatment groups (T1D-Autograft and T1D-Allograft) increased ( $6.3 \pm 1.5$  islet and  $6.0 \pm 1.0$  islet, respectively) and there were no statistically significant differences between the two groups. This showed that both autograft and allograft cell transplantation could regenerate pancreatic islet numbers. However, the size of the pancreatic islets did not increase after transplantation (Fig. 6A, B)




**Figure 6.** Comparison of the size and number of pancreatic islets in diabetic mice following mADSC treatment.

Although the size of the pancreatic islets did not increase, immunohistochemistry (IHC) staining results showed that pancreatic islets recovered their ability to secrete insulin. About 70% of cells in pancreatic islets are beta cells which are able to secrete insulin in response to blood glucose levels in mice. In islets of Normal-Placebo group, the positive staining of insulin was very clear, in contrast to the T1D-Placebo group. In the T1D-Autograft and T1D-Allograft groups, the injured pancreases were regenerated after ADSC transplantation. At day 56 post transplantation, insulin-positive cells appeared in the pancreases of treated mice. These results were confirmed by results of blood insulin levels in ELISA assays. The level of insulin that exist in the peripheral blood may reflect the regeneration of the pancreas after treatment. Insulin levels were found to be significantly different among the experimental groups (Fig. 7,8).



**Figure 7. Immunofluorescent staining of pancreatic islets for insulin.**

The results showed that the insulin concentration in treated mice (T1D-Autograft and T1D-Allograft) significantly increased compared to T1D-Placebo group ( $1.28 \pm 0.02 \mu\text{g/L}$  and  $1.58 \pm 0.13 \mu\text{g/L}$  vs.  $0.97 \pm 0.07 \mu\text{g/L}$ , respectively) (Fig. 8). These results showed that autograft and allograft ADSC transplantation helped to rehabilitate pancreas function in our diabetic mouse model.



**Figure 8.** Insulin concentration in peripheral blood of experimental groups.

## Discussion

Treatment with ADSCs has been shown to have the potential to recover pancreas. In previous studies, ADSCs have served as an ideal candidate for T1D treatment worldwide (Kono et al., 2014; Lin et al., 2009; Rahavi et al., 2015; Yaochite et al., 2015). ADSCs have been used in clinical trials based on their ability to differentiate into beta cells and stimulate regeneration of beta cells in the damaged pancreas (El-Badawy and El-Badri, 2016). Fat tissue is an abundant source of MSCs. Among the different kinds of MSCs, the proliferation of stem cells derived from adipose tissue has been demonstrated to be better than cells from bone marrow tissue (Semon et al., 2014). Simultaneously, ADSCs has a greater potential for angiogenesis (Dao et al., 2016; Vu et al., 2014).

In this study, we have successfully cultured ADSCs according to a protocol from our laboratory (Vu et al., 2015). These cells exhibited the mesenchymal stem cell phenotypes. Indeed, they could spread over the surface of plastic flasks with fibroblast shape; successfully differentiated into adipocytes and osteoblasts. According to Dominici et al. (2006), the some markers for mesenchymal stem cells should be done, however we applied the previous protocol without modifications (Vu et al., 2015), these cells only were checked their shapes and differentiation potential.

In the next experiments, ADSCs were used to treat T1D mice in two platforms included autologous and allogenic transplantations. The results showed that ADSC transplantation successfully recovered number of islets, recovered pancreas functions of diabetic mellitus mouse, and reduced blood glucose levels. More importantly, there were not significantly different about efficacy of autologous and allogenic transplantation of ADSCs in diabetes mellitus treatment. This result showed that effects of ADSCs did not dependent on autologous ADSCs or allogenic ADSCs.

Indeed, ADSCs as well as MSCs exhibited the low immunogenicity (Gu et al., 2015; Yaochite et al., 2015). Therefore, both autologous and allogenic ADSCs can exist in the models. Moreover, the effects of ADSCs can relate to some signals or growth factors that secreted by ADSCs. The secreted product of ADSCs involved in immune modulation, conditioned inflammatory process, protected the remaining beta cells and stimulated the proliferation of beta cells (Fiorina et al., 2011; Kono et al., 2014). Previous studies showed that the paracrine effects of ADSCs played important roles in controlling blood glucose in animal diabetic models. ADSCs could produce inflammatory cytokines and contribute to avoid stress due to high blood glucose (Rahavi et al., 2015).

## Conclusion

In this study, we have successfully isolated and cultured mouse adipose derived stem cells which exhibit characteristics of MSCs, such as the ability to differentiate into osteoblasts and adipocytes. After transplantation of  $2 \times 10^6$  mADSCs, mice showed recovery of pancreatic islets and improvement of insulin production of beta cells, thereby reducing blood glucose levels in mice. There was no difference between autologous (autograft) and allogeneic (allograft) transplantations of ADSCs. After treatment, blood glucose levels decreased and were stably maintained from day 35 to day 56, and mortality rate was decreased to 0%. Overall, our study shows that adipose tissue stem cells can reduce blood glucose levels and reactivate a functional part of the pancreas. Both autologous and allogeneic ADSCs are efficacious for treatment of T1D.

---

## Abbreviations

MSCs: Mesenchymal stem cells

STZ: Streptozotocin

T1D: Type 1 diabetes mellitus disease

ADSCs: Adipose-derived stem cells

## **Acknowledgment**

This research was funded by Ministry of Science and Technology via project Grant No. DTDL.2012-G/23.

## **Author Contribution**

Anh NTB, Oanh TKN, Ngoc KP isolated, confirmed and cultured ADSCs; Oanh TKN, Phuc VP analysed data, wrote and revised the manuscript; Anh NTB, Cong LTN and Dung TN produced the type 1 diabetes mellitus models by STZ; autograft and allograft transplantation of ADSCs; monitored blood glucose, glucose. Insulin tolerance assay and pancreas structure evaluation were carried out by BNT Anh; Loan TTD and Cong LTN.

## References

- Bunnell, B.A., Flaat, M., Gagliardi, C., Patel, B., and Ripoll, C. (2008). Adipose-derived stem cells: isolation, expansion and differentiation. *Methods (San Diego, Calif)* 45, 115-120.
- Dao, T.T.-T., Vu, N.B., Phi, L.T., Le, H.T.-N., Phan, N.K., and Van Pham, P. (2016). Human adipose-derived mesenchymal stem cell could participate in angiogenesis in a mouse model of acute hindlimb ischemia. *Biomedical Research and Therapy* 3, 770-779.
- El-Badawy, A., and El-Badri, N. (2016). Clinical Efficacy of Stem Cell Therapy for Diabetes Mellitus: A Meta-Analysis. *PLoS ONE* 11, e0151938.
- Fiorina, P., Voltarelli, J., and Zavazava, N. (2011). Immunological applications of stem cells in type 1 diabetes. *Endocrine reviews* 32, 725-754.
- Gu, L.H., Zhang, T.T., Li, Y., Yan, H.J., Qi, H., and Li, F.R. (2015). Immunogenicity of allogeneic mesenchymal stem cells transplanted via different routes in diabetic rats. *Cell Mol Immunol* 12, 444-455.
- Halban, P.A., Polonsky, K.S., Bowden, D.W., Hawkins, M.A., Ling, C., Mather, K.J., Powers, A.C., Rhodes, C.J., Sussel, L., and Weir, G.C. (2014). beta-cell failure in type 2 diabetes: postulated mechanisms and prospects for prevention and treatment. *Diabetes care* 37, 1751-1758.
- Karaoz, E., Okcu, A., Unal, Z.S., Subasi, C., Saglam, O., and Duruksu, G. (2013). Adipose tissue-derived mesenchymal stromal cells efficiently differentiate into insulin-producing cells in pancreatic islet microenvironment both in vitro and in vivo. *Cytotherapy* 15, 557-570.
- Katuchova, J., Harvanova, D., Spakova, T., Kalanin, R., Farkas, D., Durny, P., Rosocha, J., Radonak, J., Petrovic, D., Siniscalco, D., et al. (2015). Mesenchymal stem cells in the treatment of type 1 diabetes mellitus. *Endocrine pathology* 26, 95-103.
- Kono, T.M., Sims, E.K., Moss, D.R., Yamamoto, W., Ahn, G., Diamond, J., Tong, X., Day, K.H., Territo, P.R., Hanenberg, H., et al. (2014). Human adipose-derived stromal/stem cells protect against STZ-induced hyperglycemia: analysis of hASC-derived paracrine effectors. *Stem Cells* 32, 1831-1842.
- Kwon, D.S., Gao, X., Liu, Y.B., Dulchavsky, D.S., Danyluk, A.L., Bansal, M., Chopp, M., McIntosh, K., Arbab, A.S., Dulchavsky, S.A., et al. (2008). Treatment with bone marrow-derived stromal cells accelerates wound healing in diabetic rats. *International wound journal* 5, 453-463.
- Lin, G., Wang, G., Liu, G., Yang, L.J., Chang, L.J., Lue, T.F., and Lin, C.S. (2009). Treatment of type 1 diabetes with adipose tissue-derived stem cells expressing pancreatic duodenal homeobox 1. *Stem cells and development* 18, 1399-1406.
- Lin, H.P., Chan, T.M., Fu, R.H., Chuu, C.P., Chiu, S.C., Tseng, Y.H., Liu, S.P., Lai, K.C., Shih, M.C., Lin, Z.S., et al. (2015). Applicability of adipose-derived stem cells in type 1 diabetes mellitus. *Cell transplantation* 24, 521-532.
- Organization, W.H. (2016). Global report on diabetes.
- Paek, H.J., Kim, C., and Williams, S.K. (2014). Adipose stem cell-based regenerative medicine for reversal of diabetic hyperglycemia. *World Journal of Diabetes* 5, 235-243.

- Rahavi, H., Hashemi, S.M., Soleimani, M., Mohammadi, J., and Tajik, N. (2015). Adipose tissue-derived mesenchymal stem cells exert in vitro immunomodulatory and beta cell protective functions in streptozotocin-induced diabetic mice model. *J Diabetes Res* 2015, 878535.
- Semon, J.A., Maness, C., Zhang, X., Sharkey, S.A., Beuttler, M.M., Shah, F.S., Pandey, A.C., Gimble, J.M., Zhang, S., Scruggs, B.A., et al. (2014). Comparison of human adult stem cells from adipose tissue and bone marrow in the treatment of experimental autoimmune encephalomyelitis. *Stem cell research & therapy* 5, 2.
- Timper, K., Seboek, D., Eberhardt, M., Linscheid, P., Christ-Crain, M., Keller, U., Muller, B., and Zulewski, H. (2006). Human adipose tissue-derived mesenchymal stem cells differentiate into insulin, somatostatin, and glucagon expressing cells. *Biochemical and biophysical research communications* 341, 1135-1140.
- Vu, N.B., Bui, A.N.-T., Ngoc-Le Trinh, V., Phi, L.T., Phan, N.K., and Van Pham, P. (2014). A comparison of umbilical cord blood-derived endothelial progenitor and mononuclear cell transplantation for the treatment of acute hindlimb ischemia.
- Vu, N.B., Van Trinh, N.L., Phi, L.T., Vo, T.L.H., Dao, T.T.T., Phan, N.K., Van Ta, T., and Van Pham, P. (2015). An evaluation of the safety of adipose-derived stem cells. *Biomedical Research and Therapy* 2, 22.
- Yaochite, J.N., Caliri-Oliveira, C., de Souza, L.E., Neto, L.S., Palma, P.V., Covas, D.T., Malmegrim, K.C., Voltarelli, J.C., and Donadi, E.A. (2015). Therapeutic efficacy and biodistribution of allogeneic mesenchymal stem cells delivered by intrasplenic and intrapancreatic routes in streptozotocin-induced diabetic mice. *Stem cell research & therapy* 6, 31.
- Zhou, F., Hui, Y., Xu, Y., Lei, H., Yang, B., Guan, R., Gao, Z., Xin, Z., and Hou, J. (2016). Effects of adipose-derived stem cells plus insulin on erectile function in streptozotocin-induced diabetic rats. *International Urology and Nephrology* 48, 657-669.
- Zuk, P. (2013). Adipose-Derived Stem Cells in Tissue Regeneration: A Review. *ISRN Stem Cells* 2013, 35.